




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Outcomes of 3,737 COVID-19 patients treated with hydroxychloroquine/azithromycin and other regimens in Marseille, France: A retrospective analysis

Jean-Christophe Lagier^{a,b,1}, Matthieu Million^{a,b,1}, Philippe Gautret^{a,c}, Philippe Colson^{a,b}, Sébastien Cortaredona^{a,c}, Audrey Giraud-Gatineau^{a,c,d,e}, Stéphane Honoré^{f,g}, Jean-Yves Gaubert^h, Pierre-Edouard Fournier^{a,c}, Hervé Tissot-Dupont^{a,b}, Eric Chabrière^{a,b}, Andreas Stein^{a,b}, Jean-Claude Deharoⁱ, Florence Fenollar^{a,c}, Jean-Marc Rolain^{a,b}, Yolande Obadia^a, Alexis Jacquier^j, Bernard La Scola^{a,b}, Philippe Brouqui^{a,b}, Michel Drancourt^{a,b}, Philippe Parola^{a,c}, Didier Raoult^{a,b,*}, IHU COVID-19 Task force

^a IHU-Méditerranée Infection, Marseille, France

^b Aix Marseille University, IRD, AP-HM, MEPHI, Marseille, France

^c Aix Marseille University, IRD, AP-HM, SSA, VITROME, Marseille, France

^d Centre d'Epidémiologie et de Santé Publique des Armées (CESPA), Marseille, France

^e AP-HM, Marseille, France

^f Aix Marseille University, Laboratoire de Pharmacie Clinique, Marseille, France

^g AP-HM, Hôpital Timone, Service Pharmacie, Marseille, France

^h Department of Radiology and Cardiovascular Imaging, Aix Marseille Univ, LIIE, Marseille, France

ⁱ AP-HM, Aix Marseille University, Hôpital Timone, Cardiologie, Rythmologie, Marseille, France

^j Department of Radiology and Cardiovascular Imaging, Aix-Marseille Univ., UMR 7339, CNRS, CRMBM-CEMEREM (Centre de Résonance Magnétique Biologique et Médicale-Centre d'Exploration Métaboliques Par Résonance Magnétique), Marseille, France

ARTICLE INFO

Keywords:

SARS-CoV-2

COVID-19

Hydroxychloroquine

Azithromycin

ABSTRACT

Background: In our institute in Marseille, France, we initiated early and massive screening for coronavirus disease 2019 (COVID-19). Hospitalization and early treatment with hydroxychloroquine and azithromycin (HCQ-AZ) was proposed for the positive cases.

Methods: We retrospectively report the clinical management of 3,737 screened patients, including 3,119 (83.5%) treated with HCQ-AZ (200 mg of oral HCQ, three times daily for ten days and 500 mg of oral AZ on day 1 followed by 250 mg daily for the next four days, respectively) for at least three days and 618 (16.5%) patients treated with other regimen ("others"). Outcomes were death, transfer to the intensive care unit (ICU), ≥ 10 days of hospitalization and viral shedding.

Results: The patients' mean age was 45 (sd 17) years, 45% were male, and the case fatality rate was 0.9%. We performed 2,065 low-dose computed tomography (CT) scans highlighting lung lesions in 592 of the 991 (59.7%) patients with minimal clinical symptoms (NEWS score = 0). A discrepancy between spontaneous dyspnoea, hypoxemia and lung lesions was observed. Clinical factors (age, comorbidities, NEWS-2 score), biological factors (lymphocytopenia; eosinopenia; decrease in blood zinc; and increase in D-dimers, lactate dehydrogenase, creatinine phosphokinase, troponin and C-reactive protein) and moderate and severe lesions detected in low-dose CT scans were associated with poor clinical outcome. Treatment with HCQ-AZ was associated with a decreased risk of transfer to ICU or death (Hazard ratio (HR) 0.18 0.11–0.27), decreased risk of hospitalization ≥ 10 days (odds ratios 95% CI 0.38 0.27–0.54) and shorter duration of viral shedding (time to negative PCR: HR 1.29 1.17–1.42). QTc prolongation (>60 ms) was observed in 25 patients (0.67%) leading to the cessation of treatment in 12 cases including 3 cases with QTc > 500 ms. No cases of *torsade de pointe* or sudden death were observed.

* Corresponding author. IHU - Méditerranée Infection, 19-21 boulevard Jean, Moulin, 13005, Marseille, France.

E-mail address: didier.raoult@gmail.com (D. Raoult).

¹ Equal contribution.

<https://doi.org/10.1016/j.tmaid.2020.101791>

Received 27 May 2020; Received in revised form 12 June 2020; Accepted 14 June 2020

Available online 25 June 2020

1477-8939/© 2020 The Authors.

Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Conclusion: Although this is a retrospective analysis, results suggest that early diagnosis, early isolation and early treatment of COVID-19 patients, with at least 3 days of HCQ-AZ lead to a significantly better clinical outcome and a faster viral load reduction than other treatments.

1. Introduction

In December 2019, a novel coronavirus emerged in the central Chinese province of Hubei, causing an outbreak of pneumonia [1]. As of June 11th, 2020, more than 7 million persons were infected with SARS-CoV-2, and more than 400,000 have died [2]. Management of this infection was heterogeneous across countries regarding i) indications for virological testing of patients and asymptomatic contacts, ii) indications for low-dose computed tomography (LDCT), and iii) therapeutic options and follow-up. Based on preliminary data from Chinese physicians [3,4], in Marseille, France, we designed a strategy including early massive screening by PCR, LDCT of the chest for positive patients, and early treatment with hydroxychloroquine (HCQ) to which we rapidly added azithromycin (AZ) after we found that the combination had a synergistic effect against the virus *in vitro* [5] and *in vivo* [6–8]. This led our institute to face a dramatic increase in workload but allowed us to generate real-life data allowing us to comprehensively describe the disease and management at our institute, despite the inherent limitations of such an observational study (Table 1).

Indeed, among the candidate treatments, only three main drugs (remdesivir, lopinavir-ritonavir and HCQ) have been tested in large comparative studies [11–13]. Lopinavir-ritonavir and remdesivir have not clearly demonstrated efficacy but are associated with many adverse events [11,12,14]. HCQ has demonstrated its efficacy in reducing viral shedding persistence [6] and improving clinical status in observational or randomized clinical trials [13,15,16]. In addition, we performed a recent meta-analysis of 20 available reports, including 105,040 patients demonstrating that, in clinical studies, chloroquine and its derivatives improve clinical and biological outcomes and reduce mortality by a factor 3 in coronavirus disease 2019 (COVID-19) patients [10]. In addition, we recently reported a very low mortality rate in a retrospective analysis of more than 1,000 patients early treated with a combination of HCQ-AZ, with a very low mild adverse event rate (2.3%) [8]. Conversely, in a recent observational study, patients treated with HCQ showed no difference regarding risk of death or intubation compared with patients under other treatments [17]. However, the patients included in the group receiving HCQ had more severe disease and had more comorbidities than those who did not receive the drug [17].

Here, we report on more than 3,700 cases treated in our institute, including those previously reported [7,8], to give a comprehensive analysis of our strategy. Outcomes were death, transfer to intensive care unit (ICU), hospitalization stay ≥ 10 days and viral shedding persistence.

2. Materials and methods

2.1. Patients and study design

The study was conducted in the Institut Hospitalo-Universitaire (IHU) *Méditerranée Infection* (<https://www.mediterranee-infection.com/>), Assistance Publique-Hôpitaux de Marseille (AP-HM), Southern France. As previously described, we performed early massive PCR screening both for patients suspected of having COVID-19 and for contacts of confirmed cases [8]. We proposed standardized treatment and follow-up for all individuals >18 years of age with PCR-documented SARS-CoV-2 RNA from a nasopharyngeal sample. Data were collated from all patients from March 3rd to April 27th and were analysed retrospectively.

Table 1

Key numbers of activities at IHU Méditerranée Infection (2020, February 27th – 2020 May 12th).

Patients tested for SARS-CoV-2	31,003 individuals including 1,277 health care workers
Patients hospitalized in day-care hospital	3,525
Patients hospitalized in infectious diseases units	705
Serology SARS-CoV-2	6,000 samples tested including 643 samples from health care workers
Culture	4,786 samples inoculated 1,908 SARS-CoV-2 strains isolated
Genome	466 genomes sequenced and analysed
Low-dose CT scan	2,218 performed
Electrocardiograms	7,800 performed
Serum drug dosages	1,939 hydroxychloroquine dosages

2.2. Clinical, biological and radiological data and follow-up

Demographics (age, sex), chronic conditions (cancer, diabetes mellitus, chronic heart disease, hypertension, chronic respiratory disease and obesity) and concomitant medications were documented. Symptoms, including rhinitis, anosmia, ageusia, fever, cough, dyspnoea and chest pain, were systematically recorded. Severity was assessed using the National Early Warning Score adapted to COVID-19 patients (NEWS-2) at admission and during follow-up [18]. Three categories of clinical worsening were defined: low score (NEWS-2 = 0–4), medium score (NEWS-2 = 5–6), and high score (NEWS-2 ≥ 7).

We recorded lymphocyte, eosinophil and platelet counts; fibrinogen; D-dimer and other coagulation factors; electrolytes; zinc; lactate dehydrogenase (LDH); creatine phosphokinase (CPK); C-reactive protein; and HCQ serum dosage [19]. Viral load was analysed by qPCR from nasopharyngeal swabs [8] at admission and during the follow-up, and an indirect immunofluorescence quantitative assay was used to assess the serological status against SARS-CoV-2 [20]. Viral culture was attempted for PCR-positive patients [5]. A LDCT was proposed for all patients when possible. Radiological lung lesions were classified into three categories: minimal, intermediate and severe involvement [8].

2.3. COVID-19 management and outcomes

The treatment consisted of the combination of HCQ (200 mg of oral HCQ, three times daily for ten days) and AZ (500 mg on day 1 followed by 250 mg daily for the next four days). This regimen was proposed as standard care for all patients without contraindications to these drugs [8]. Patients were informed of the off-label character of the prescription of HCQ and AZ prior to receiving treatment. Treatment was initiated among inpatients in our day-care hospital (i.e. here are patient kept just during the day) or in our infectious disease hospitalization units. All patients underwent electrolyte analysis and an electrocardiogram (EKG) with corrected QT measurement (Bazett's formula) before starting treatment [8]. EKGs with any abnormalities were systematically referred to a cardiologist for further assessment. In addition, broad-spectrum antibiotics (ceftriaxone or ertapenem) were included in the regimen for patients with pneumonia and/or NEWS scores ≥ 5 . Standard care included systematic oxygen supplementation and preventive anticoagulation when necessary.

As it is common practice to assess the clinical evolution at 72 h for pneumonia [21], we selected this time-point to evaluate the clinical efficacy of HCQ-AZ [8]. Therefore, we defined two groups of patients: i) those receiving HCQ-AZ for at least three days and ii) the others




Fig. 1. Flowchart summarizing our study design.

comprising treatment with HCQ alone, AZ alone, HCQ-AZ for less than 3 days before defined clinical outcome, and those receiving neither HCQ either AZ.

Poor clinical outcome was defined as one of the following outcomes (transfer to ICU, death, hospitalization lasting ≥ 10 days), while others were considered as having a good clinical outcome.

2.4. Statistical methods

We used the Wilcoxon Mann-Whitney U test, χ^2 test, or Fisher's exact test to compare differences between groups of patients where appropriate. We performed multiple correspondence analysis (MCA) to investigate the associations between clinical data, biological data, radiological data, poor clinical outcome and the treatment received (HCQ-AZ for at least three days, other treatments). Visual observations of Kaplan-Meier curves and log-rank tests were used for survival analyses. Multivariate logistic regression and the Cox proportional hazard model were used to identify independent predictors of each outcome. Considering that death was a main outcome and that only 35 patients died in our cohort (0.9%), the number of covariates to be included in multivariate analyses was *a priori* limited to three variables: previous health status (modified Charlson combined comorbidity index) [22], severity of the disease (NEWS-2 score) and treatment (HCQ + AZ ≥ 3 days). Association between treatment (HCQ + AZ ≥ 3 days) and death was estimated by Cox regression models using three different methods. In the primary analysis, a multivariable Cox regression adjusted on the combined comorbidity index and the NEWS score was performed. We conducted a secondary analysis that used propensity-score matching. The propensity score was calculated using multivariable logistic regression on the combined comorbidity index and the NEWS score. Each patient of the "other treatment" group was matched to a patient selected of the "HCQ-AZ ≥ 3 days" group using the 1:1 nearest-neighbour propensity score matching method to create a matched sample. The third analysis used inverse probability weighting (<https://cdn1.sph.harvard.edu/wp-content/uploads/sites/343/2013/03/msm-web.pdf>). Association between treatment and death was estimated using stratified and weighted Cox regression. A two-sided p -value

of less than 0.05 was considered statistically significant. MCA was performed using R Statistical Software and the FactoMineR package. All other analyses were carried out using SAS 9.4 statistical software (SAS Institute, Cary, NC).

2.5. Ethics statement

Data presented herein were collected retrospectively from the routine care setting using the electronic health recording system of the hospital. This non-interventional retrospective study was approved by our institutional review board committee (Méditerranée Infection N°: 2020-021). In France, at the time the study was conducted, HCQ for COVID-19 treatment was approved off-label for hospital delivery only. As previously reported [8], for all patients, the prescription of HCQ-AZ was made during either complete hospitalization or at day-care hospital by one of the physicians, after collegial decision based on the most recent scientific data available and after assessment of the benefit/harm ratio of the treatment. According to European General Data Protection Regulation No 2016/679, patients were informed of the potential use of their medical data and that they could refuse the use of their data. The analysis of collected data followed the reference methodology MR-004 registered on N° MR 5010010520 in the AP-HM register.

3. Results

3.1. Selection of the current cohort

From March 3rd to April 27th, we performed 101,522 SARS-CoV-2 PCR tests among 65,993 people (including more than 25,302 sampled in the IHU). Among them, 6,831 patients tested positive (10.4%). Of these, 3,024 patients (comparable in age and sex) were excluded: 1,399 whose samples were sent to our laboratory but who were followed up outside Marseille and 1,363 patients who were managed in Marseille, outside IHU. Among the patients diagnosed and treated in the IHU, 3,737 were analysed in this study after the exclusion of 332 younger than 18 years of age (described elsewhere).

Table 2
Baseline characteristics of the patients according to treatment.

	All		HCQ-AZ ≥ 3 days		Other treatments		HCQ-AZ < 3 days		HCQ		AZ		No HCQ, No AZ	
	n = 3,737		(n = 3,119 83.5%)		(n = 618 16.5%)		(n = 218 5.8%)		(n = 101 2.7%)		(n = 137 3.7%)		(n = 162 4.3%)	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Age														
Age 18-44	1874	50.2	1649	52.8	225	36.4*	76	34.9*	46	45.5	24	17.5*	79	48.8*
Age 45-54	804	21.5	671	21.5	133	21.5	50	22.9	29	28.7	22	16.1	32	19.7
Age 55-64	606	16.2	503	16.1	103	16.7	38	17.4	17	16.8	25	18.2	23	14.2
Age 65-74	241	6.5	183	5.9	58	9.4	21	9.6	4	4	20	14.6	13	8
Age >74	212	5.7	113	3.6	99	16	33	15.1	5	4.9	46	33.6	15	9.3
Sex														
Men	1704	45.6	1416	45.4	288	46.6	105	48.2	47	46.5	64	46.7	72	44.4
Chronic condition(s)														
Cancer disease	129	3.5	83	2.7	46	7.4*	20	9.2*	3	3	16	11.7*	7	4.3
Diabetes	312	8.4	235	7.5	77	12.5*	23	10.5	4	4	37	27.0*	13	8
Chronic heart diseases	219	5.9	125	4	94	15.2*	23	10.5*	7	6.9	46	33.6*	18	11.1*
Hypertension	561	15	410	13.1	151	24.4*	50	22.9*	13	12.9	57	41.6*	31	19.1*
Chronic respiratory diseases	338	9	267	8.6	71	11.5*	21	9.6	9	8.9	25	18.2*	16	9.9
Obesity	418	11.2	345	11.1	73	11.8	25	11.5	5	4.9	28	20.4*	15	9.3
Symptom(s) declared by patient ^a	3397	90.9	2862	91.8	535	86.6*	202	92.7	86	85.1*	116	84.7*	131	80.9*
Fever	574	15.6	468	15.1	106	18.6*	42	22.5*	25	25.0*	20	16.3	19	11.9
Cough	1846	50.2	1578	50.8	268	47.1*	88	47.1	56	56	51	41.5*	73	45.9
Rhinitis	1202	32.7	1065	34.3	137	24.1*	46	24.6*	28	28	21	17.1*	42	26.4*
Anosmia	1442	39.2	1277	41.1	165	29.0*	59	31.5*	28	28.0*	25	20.3*	53	33.3
Ageusia	1389	37.8	1213	39	176	30.9*	61	32.6	33	33	27	21.9*	55	34.6
Dyspnea	1038	28.2	901	29	137	24.1*	65	34.8	16	16.0*	26	21.1	30	18.9*
Thoracic pain	811	22.1	745	24	66	11.6*	27	14.4*	12	12.0*	7	5.7*	20	12.6*
NEWS score														
0-4	3420	91.5	2925	93.8	495	80.1*	165	75.7*	94	93.1	91	66.4*	145	89.5*
5-6	172	4.6	114	3.7	58	9.4	22	10.1	5	4.9	25	18.2	6	3.7
>6	145	3.9	80	2.6	65	10.5	31	14.2	2	2	21	15.3	11	6.8
Pulmonary CT-scanner ^b														
Normal	616	29.8	540	31.6	76	21.4*	28	19.3*	9	23.1	19	18.3*	20	29.9
Minimal	928	44.9	780	45.6	148	41.7	55	37.9	25	64.1	38	36.5	30	44.8
Intermediate	414	20.1	329	19.2	85	23.9	36	24.8	4	10.2	30	28.8	15	22.4
Severe	107	5.2	61	3.6	46	13	26	17.9	1	2.6	17	16.3	2	3

*: p < 0.05 (Fisher's exact test). Reference group is "HCQ-AZ ≥ 3 days", ^aData available for 3676 patients. One patient may present several symptoms. ^bData available for 1710 patient in the "HCQ-AZ ≥ 3 days" group, 145 in the "HCQ-AZ < 3 days" group, 39 in the "HCQ only" group, 104 in the "AZ only" group and 67 in the "other treatments" group.

3.2. Overall characteristics of patients

Among the 3,737 included patients, 3,284 (87.9%) were younger than 65 years, and 453 were older (12.1%), with a mean age of 45.3 years (standard deviation (sd), 16.8). A total of 1,704 patients (45.6%) were male. Regarding therapeutic management, 3,119 (83.5%) patients received at least a 3-day course of HCQ-AZ. Among the 618 others, 218 received a shorter course of HCQ-AZ (35.3%), 137 received AZ alone (22.2%), 101 received HCQ alone (16.3%) and 162 did not receive either drug (26.2%) (Fig. 1). The baseline characteristics of patients according to treatment groups are summarized in Table 2. We paid a rigorous attention to avoiding HCQ-AZ in patients with cardiac diseases, abnormal EKG, dyskalemia or current use of other interacting medications (Table 3).

Overall, 673 patients (18%) were hospitalized in our infectious disease units, and 3,064 patients were followed in our day-care hospital (Fig. 1). Most of the patients (3,507, 93.8%) had a good clinical outcome, while 230 (6.2%) had a poor clinical outcome, including 67 who were transferred to ICU (1.8%), 35 who died (0.9%) and 197 with a hospital stay ≥10 days (5.3%) (Table 4).

The multiple correspondence analysis (MCA) (Fig. 2) immediately allowed for the identification of a number of groups. Most patients with a good clinical outcome are grouped with young age and centred on HCQ-AZ treatment. The patients with a poor clinical outcome are all grouped with older age, with some biological criteria (lymphocytopenia, thrombocytopenia, eosinopenia, low zinc level and increased D-dimers

and troponin) and with other treatments. Finally, the two modes of clinical presentation are highlighted: upper respiratory tract infection (URTI) symptoms with ageusia, anosmia, rhinitis, and thoracic pain, and lower respiratory tract infection symptoms (LRTIs) with dyspnoea, cough and fever (Fig. 2).

3.3. Clinical characteristics

Underlying conditions, comedications and clinical symptoms are comprehensively described in Table S1. The prevalence of poor clinical outcome significantly increased with age, comorbidities, several comedications and male sex.

Most of the patients had a NEWS-2 score ranging from 0 to 4 (3,420, 91.5%) at admission. Cough was the most frequent symptom (50.2%), followed by anosmia (39.2%), ageusia (37.8%), rhinitis (32.7%), dyspnoea (28.2%) and thoracic pain (22.1%). A total of 15.6% of patients were febrile, and 9.1% were asymptomatic. Interestingly, anosmia, ageusia and chest pain were significantly more frequent in patients under 65 years (42.9% vs 11.4%, 40.9% vs. 14.5% and 24.3% vs. 4.9%, respectively).

Symptoms suggestive of URTI, including rhinitis, anosmia and ageusia, were significantly more common in patients with a good clinical outcome (33.8% vs. 15.6%, 40.9% vs. 11.9% and 39.3% vs. 14.2%, respectively). The symptoms suggestive of LRTI, including fever and dyspnoea, were significantly more frequent in patients with poor clinical outcome (32.1% vs. 14.6% and 40.8% vs. 27.4%, respectively).

Table 5
Age stratified multivariable analyses adjusted on comorbidities and severity of the disease addressing associations between treatment (HCQ-AZ \geq 3 days) and clinical outcomes/viral shedding clearance (n = 3,737).

Cox proportional hazard models ^a	HCQ-AZ \geq 3 days n event/n total (%)	Other treatment n event/n total (%)	Hazard ratio 95% confidence interval (ref. Other treatment)	p-value
Mortality^b	16/503 (3.2%)	19/199 (9.6%)		
Multivariable Cox regression on unmatched sample (n = 702)			0.49 0.25–0.97	0.0406
Stratified Cox regression on matched sample (n = 398) ^c			0.41 0.17–0.99	0.0482
Weighted Cox regression on unmatched sample (n = 702) ^c			0.49 0.31–0.79	0.0030
ICU transfer^d				
Patients \geq 31 years (n = 2,856)	25/2,355 (1.1%)	42/501 (8.4%)	0.19 0.11–0.33	<0.0001
Patients between 31 and 59 years (n = 2,180)	10/1,862 (0.5%)	23/318 (7.2%)	0.13 0.05–0.31	<0.0001
Patients aged \geq 60 years (n = 676)	15/493 (3.0%)	19/183 (10.4%)	0.17 0.07–0.38	0.0003
Death and/or ICU transfer^e				
Patients \geq 31 years (n = 2,882)	35/2,365 (1.5%)	58/517 (11.2%)	0.18 0.11–0.27	<0.0001
Patients aged \geq 60 years (n = 702)	25/503 (5.0%)	35/199 (17.6%)	0.30 0.18–0.51	<0.0001
Viral shedding persistence \geq 10 days^f				
All patients (n = 3,737)	10.6%	20.6%	1.29 1.17–1.42	<0.0001
Patients aged < 60 years (n = 3,035)	10.0%	17.4%	1.23 1.10–1.38	0.0003
Patients aged \geq 60 years (n = 702)	13.4%	27.2%	1.44 1.19–1.73	0.0002
Logistic regression^g	HCQ-AZ \geq 3 days n event/n total (%)	Other treatment n event/n total (%)	Odds ratio 95% confidence interval	p-value
Hospitalization \geq 10 days				
All patients (n = 3737)	109/3,119 (3.5%)	88/618 (14.2%)	0.38 0.27–0.54	<.0001
Death and/or ICU transfer/hospitalization \geq 10 days				
All patients (n = 3737)	121/3,119 (3.9%)	109/618 (17.6%)	0.30 0.22–0.42	<.0001

^a Models were adjusted for the combined comorbidity index and the severity of the disease (NEWS-2 score).

^b Mortality was evaluated among patients aged 60 years old and older (n = 702) because the youngest patient who died was 60 years old.

^c These two models based on propensity score methods were performed only for mortality (see methods).

^d ICU transfer was evaluated among patients aged 31 years and older (n = 2,856) because the youngest patient who was transferred to the ICU was 31 years old. Patients who died without ICU transfer were excluded (n = 26).

^e Death and/or ICU transfer was evaluated among patients aged 31 years and older (n = 2,856) because the youngest patient who was transferred to the ICU was 31 years old.

^f Proportion of patients with non-negative PCR within 10 days following inclusion (Kaplan-Meier estimates, see Fig. 3). Some patients did not have a PCR

testing at day 10 and were still considered positive if previous sample was positive (event was defined as first negative PCR during follow-up).

among patients treated with HCQ-AZ \geq 3days. Whereas no death was observed in patients <60 years old in our study, the proportion of deaths under 60 years was 3.5, 4.3, 9.8 and 19% in Italy, in grand Est region (France), in Ile de France region (France) and in China, respectively (Table 6) [23].

3.8. Virological outcome

Kaplan-Meier estimates show that the proportion of patients with positive PCR 10 days after inclusion was significantly lower among patients treated with HCQ-AZ (10.6%; (95% CI: 8.1%–13.4%)) than among those who received other treatments (20.6%; (95% CI: 14.7%–27.2%; p < 0.05) (Fig. 3, Table 5). In a multivariate Cox regression adjusted for combined comorbidity index and disease severity at admission (NEWS-2 score), HCQ-AZ treatment remained significantly associated with viral shedding clearance (HR = 1.29: 1.17–1.42, p < 0.0001) (Table 5). We inoculated samples obtained from 130 patients with positive PCR at day 10. Among them, only 16 had a positive culture at day 10 (12.3%).

4. Discussion

This work highlights that it is hazardous to make strategic decisions *a priori* regarding the management of a new disease when no reliable information about this disease is available. Political and public health decisions in this context should be regularly adapted to observations collected in other countries when available [24]. The decision of the government of France to recommend staying at home (lockdown) without testing while waiting for dyspnoea was not supported by our results [25]. As with other clinicians, we have seen patients with hypoxia, including some with very low blood oxygen levels, who described themselves as feeling well and comfortable (“happy hypoxemia”) [26]. Since these patients may develop severe symptoms based on our observations, the use of inexpensive pulse oximeters (around 20€) in primary-care health settings and/or by family doctors might be considered a triage tool on which to base hospitalization referral for further investigation. We propose that the initial disease severity assessment cannot rely only on clinical examination but should also take into account oxygen saturation testing and blood sampling (haemogram, CRP, LDH, troponin, D-dimers) (Fig. 2).

We confirm here that COVID-19 has several evolutionary stages (Fig. 4). After the incubation period, the first clinical stage, including LRTI and URTI symptoms, is associated with a high viral load and the occurrence of early lung lesions on LDCT, for which it is reasonable to use a compound with antiviral activity. HCQ-AZ has demonstrated its effectiveness in reducing viral shedding [6] and preventing disease progression and death particularly when prescribed at early stages [10, 27,28]. Other antiviral compounds, including remdesivir and hyperimmune gamma globulins [29], may have antiviral activity at an early stage of the disease, although there is to date no convincing published report, comparable to that of oseltamivir at the early stage of influenza [30]. Taking into account the association between low blood zinc levels and poor clinical outcomes, zinc supplementation should be also considered, as recently reported [31]. However, the choice of the best treatment should be made according to its safety profile, which is much better for HCQ-AZ than for remdesivir (adverse events leading to cessation of treatment in 0.9% in our study vs. 12% for remdesivir [12]). Nevertheless, we were surprised by the large discrepancy on efficacy and toxicity of HCQ in recent studies compared to ours [32]. As a matter of fact, all patients reported here have been followed by the physicians authors named in our study. Altogether, we found only 0.67% of QTc prolongations and no death related to treatment. In our opinion, this excellent safety profile of HCQ-AZ in our real-life medical experience




Fig. 3. Kaplan-Meier curve of clinical outcomes/viral shedding clearance according to treatment groups (n = 3,737). HCQ: hydroxychloroquine, AZ: azithromycin, ICU: Intensive care unit, PCR: polymerase chain reaction. a: For time to negative PCR, event was defined as first negative PCR during follow-up. Accordingly, patients were still considered positive at each time point if previous sample was positive.

much better reflects the reality than registry studies such as those recently retracted from high profile medical journals [9].

The second stage includes both an immune reaction and the persistence of the virus [1]. At this stage, extreme caution should be required for patients with risk factors (particularly hypertension), severe clinical presentation (NEWS CoV ≥ 5), intermediate-to-severe lesions in LDCT and biological parameters such as lymphocytopenia, eosinopenia, elevated troponin or D-dimers higher than 0.5 µg/L. Systemic coagulation activation and thrombotic complications were probably overlooked in COVID-19 patients. In our study, the youngest person who died was 60 years old, and the death was associated with generalized thrombosis. A recent study reported that among 198 hospitalized COVID-19 patients, 39 (20%) were diagnosed with venous thromboembolism (VTE), and of these patients 25 (13%) had symptomatic VTE, despite routine thrombosis prophylaxis [33]. The third stage consists of an inflammatory stage linked to pro-inflammatory cytokine release with a high risk of transfer to ICU [34]. Moreover, the strong specific antibody response observed at this stage questions the use of hyperimmune gamma globulins [29]. The fourth stage with acute respiratory distress syndrome (ARDS) is characterised by pulmonary tissue injury and requires supportive intensive care. To date, no drug has proven effective at this stage. While most surviving patients may be definitely cured, an unknown proportion may evolve towards pulmonary fibrosis constituting the late stage of the disease, as described by Chinese physicians caring for COVID-19 patients and as previously described for severe acute respiratory syndrome (SARS) in 2003 [35]. Long-term follow-up aiming to screen for fibrosis will be the next challenge in the management of COVID-19. Our experience and suggestions regarding the various stages of COVID-19 are summarized in Fig. 4.

The strength of our study is its monocentric design with a relative homogeneity of both diagnosis procedure and standard care provided to patients, allowing us to assess the impact of different therapeutic options on the evolution of the disease, in real time. Virological diagnosis, radiological investigations and clinical assessment were conducted by single teams of trained virologists, radiologists, infectious diseases specialists and cardiologists, all directly involved in patient care. Daily staff meeting ensured assessing the reliability of the data collected and adjusting medical procedures overtime, in the context of a newly emerging disease that was totally unknown three months before we started our study. In the context of pandemic, such a study allows more flexibility than a randomized controlled trial (RCT) with stringent methodological constraints and is much more economical. It is also more reliable than big-data studies conducted by external investigators dealing with incomplete information retrieved from medical files that have led to recent retraction of papers from the two major medical journals [9]. Indeed, when inconsistencies appeared in our database of the study, the authorized person in our team was able to return to patients files to reassess the data. In addition, we were able to conduct interim analysis of our data [6–8] and ensure early release of our preliminary results to be shared with the medical community, at an early phase of the pandemic [36].

Our study has a retrospective observational design, and such characteristics may be presented as a limitation of the study [37]. Patients were not enrolled in perfectly homogeneous groups with regards to demographics, chronic conditions and clinical status at admission. Treatments were not allocated randomly but according to the clinical status of patients and contra-indications to drugs, or preference of patients with regards to therapeutic options. As we have aimed to test and treat all positive patients presenting to our institution, the patient population comprised a majority of patients with mild diseases and a minority of patients with severe disease, with the former managed at our day-care hospital and the latter as in-patients. We enrolled all patients including those who started their treatment with delay or stopped it early. Because of the crisis situation we had to face, clinical, virological and radiological data were not documented in 100% patients. However, missing data may also be a limitation of RCT. Furthermore, RCT are not

Table 6

Numbers of deaths in hospitalized COVID-19 patients and distribution by age class in Italy, China, IHU Méditerranée Infection, Marseille France, Grand Est region and Ile de France regions of France.

Age class	Italy as of March 17, 2020 (Onder, 2020) ^a	China as of February 11, 2020 (Onder, 2020) ^a	IHU All patients April 30, 2020	IHU April 30, 2020 HCQ-AZ at least 3 days	IHU April 30, 2020 Other treatments	Grand-Est region, France May 18, 2020 ^b	Ile-de-France, France May 18, 2020 ^b
All	1,624	1,023	35	16	19	3,277	6,713
0-9	0	0	0	0	0	1 (0.03%)	2 (0.03%)
10-19	0	1 (0.1%)	0	0	0	0	3 (0.04%)
20-29	0	7 (0.7%)	0	0	0	2 (0.06%)	11 (0.2%)
30-39	4 (0.2%)	18 (1.8%)	0	0	0	15 (0.5%)	45 (0.7%)
40-49	10 (0.6%)	38 (3.7%)	0	0	0	32 (1.0%)	124 (1.8%)
50-59	43 (2.6%)	130 (12.7%)	0	0	0	91 (2.8%)	470 (7.0%)
60-69	139 (8.6%)	309 (30.1%)	2 (5.7%)	1 (6.25%)	1 (5.3%)	350 (10.7%)	982 (14.6%)
70-79	578 (35.6%)	312 (30.5%)	14 (40%)	7 (43.75%)	7 (36.8%)	818 (25.0%)	1,586 (23.6%)
≥80	850 (52.3%)	208 (20.3%)	19 (54.3%)	8 (50%)	11 (57.9%)	1,968 (60.1%)	3,490 (52.0%)
<60	57 (3.5%)	194 (19.0%)	0 (0%)	0 (0%)	0 (0%)	141 (4.3%)	655 (9.8%)
≥60	1567 (96.5%)	829 (81.0%)	35 (100%)	16 (100%)	19 (100%)	3136 (95.7%)	6058 (90.2%)

^a Mortality data provided in this study are likely to be global and not only that of hospitalized patients.

^b These data are collected by Sante Publique France (<https://geodes.santepubliquefrance.fr/#view=map2&c=indicateur>); IHU (Institut Hospitalo Universitaire), Marseille, France.




Fig. 4. Evolutionary stages of SARS-CoV-2 infection, including major clinical and biological features and possible therapies.

useful in the context of an emerging pandemic when commercially available drugs known to be active *in vitro* are available for immediate treatment [38,39].

Our approach of early diagnosis and care of as many patients as possible results in much lower mortality rates than other strategies. The test-and-treat strategy adopted in Marseille also seems capable of shortening the duration of the outbreak when compared to data from

France overall by identifying infected people and reducing their viral shedding duration. In fact, more people were tested in Marseille than in most other areas, and the outbreak lasted only 9 weeks. In addition, patients under HCQ-AZ treatment for at least 3 days had a better clinical outcome, based on mortality rates among patients >60 years, less transfer to ICU and shorter length of stay at the hospital, and these patients also had a shorter duration of viral shedding than patients who did

not receive this drug combination. Finally, a global strategy for the management of the COVID-19 outbreak may help to limit both the number of cases and fatalities and guide countries where this pandemic has not yet peaked.

Author's note

Since this analysis was completed, and as of the 11th June, 2020, 6 more patients died including 1 patient treated with HCQ-AZ for at least 3 days and 5 in the other group, resulting in an overall 1.1% case fatality rate for the 3,737 patients included in our study.

Funding

This work was funded by ANR "Investissements d'avenir", Méditerranée infection 10-IAHU-03, and was also supported by Région Provence-Alpes-Côte d'Azur. This work had received financial support from the Mediterranean Infection Foundation.

Declaration of competing interests

The authors declare no competing interests. Funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript. Our group used widely available generic drugs distributed by many pharmaceutical companies.

CRediT authorship contribution statement

Jean-Christophe Lagier: Conceptualization, Investigation, Formal analysis, Writing - original draft. **Matthieu Million:** Conceptualization, Formal analysis, Writing - original draft. **Philippe Gautret:** Investigation, Formal analysis, Writing - original draft. **Philippe Colson:** Investigation. **Sébastien Cortaredona:** Formal analysis. **Audrey Giraud-Gatineau:** Investigation. **Stéphane Honoré:** Investigation. **Jean-Yves Gaubert:** Investigation. **Pierre-Edouard Fournier:** Investigation, Formal analysis. **Hervé Tisotto-Dupont:** Investigation. **Eric Chabrière:** Investigation. **Andreas Stein:** Investigation. **Jean-Claude Deharo:** Investigation. **Florence Fenollar:** Investigation. **Jean-Marc Rolain:** Investigation. **Yolande Obadia:** Formal analysis. **Alexis Jacquier:** Investigation. **Bernard La Scola:** Investigation. **Philippe Brouqui:** Investigation, Formal analysis. **Michel Drancourt:** Investigation. **Philippe Parola:** Conceptualization, Formal analysis, Writing - original draft. **Didier Raoult:** Conceptualization, Formal analysis, Writing - original draft.

Acknowledgements

We thank the 7 reviewers who helped in substantially improving and clarifying the manuscript with their many comments and suggestions.

We are thankful to:

Julien Andreani, Marion Bechet, Amin Ben Lassoued, Corinne Blasco, Céline Boschi, Laurent Boyer, Jean-Paul Casalta, Hervé Chaudet, Florian Correard, Pierre Dudouet, Grégory Dubourg, Carole Eldin, Vera Esteves-Vieira, Véronique Filosa, Léa Fuster, Marc Gainier, Lugdivine Ganci, Céline Gazin, Stephanie Gentile, Clio Grimaldier, Véronique Jacomo, Marie-Thérèse Jimeno, Elisabeth Jouve, Alexandra Kotovtchikhine, Béatrice Labric, Marc Léone, Nicolas Levy, Pierre-Yves Lévy, Léa Luciani, Cléa Melenotte, Lucille Pinault, Pierre Pinzelli, Elsa Prudent, Laurence Thomas, Joana Vitte, Nathalie Wurtz.

All medical students from Aix Marseille University, all nurses, laboratory staff, administrative, technical and security staff of *Assistance Publique-Hôpitaux de Marseille* and *IHU Méditerranée Infection*, all volunteer medical doctors, and the *Bataillon des Marins Pompiers de Marseille* for their help.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.tmaid.2020.101791>.

References

- [1] Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–62.
- [2] Johns Hopkins University. Coronavirus resource center. COVID-19 dashboard by the center for systems science and engineering (CSSE) at Johns Hopkins University (JHU). <https://coronavirus.jhu.edu/map.html>. [Accessed 13 May 2020].
- [3] Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res* 2020;30(3): 269–71.
- [4] Gao J, Tian Z, Yang X. Breakthrough: chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *Biosci Trends* 2020;14(1):72–3.
- [5] Andreani J, Le Bideau M, Duflot I, et al. In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect. *Microb Pathog* 2020:104228.
- [6] Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents* 2020:105949 [Epub ahead of print].
- [7] Gautret P, Lagier JC, Parola P, et al. Clinical and microbiological effect of a combination of hydroxychloroquine and azithromycin in 80 COVID-19 patients with at least a six-day follow up: an observational study. *Trav Med Infect Dis* 2020 Apr 11:101663 [Epub ahead of print].
- [8] Million M, Lagier JC, Gautret P, et al. Early treatment of COVID-19 patients with hydroxychloroquine and azithromycin: a retrospective analysis of 1061 cases in Marseille, France. *Trav Med Infect Dis* 2020 May 5:101738. <https://doi.org/10.1016/j.tmaid.2020.101738> [Epub ahead of print].
- [9] Ledford H, Van Noorden R. High-profile coronavirus retractions raise concerns about data oversight. *Nature* 2020 Jun 5. <https://doi.org/10.1038/d41586-020-01695>.
- [10] Million M, Gautret P, Colson P, et al. Clinical efficacy of chloroquine derivatives in COVID-19 infection : comparative meta-analysis between the big data and the real world. *New Microbes New Infect* 2020. <https://doi.org/10.1016/j.nmni.2020.100709>. in press.
- [11] Cao B, Wang Y, Wen D, et al. A trial of lopinavir-ritonavir in adults hospitalized with severe covid-19. *N Engl J Med* 2020;382(9):1787–99.
- [12] Wang Y, Zhang D, Du G, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet* 2020;395 (10236):1569–78.
- [13] Yu B, Li C, Chen P, et al. Low dose of hydroxychloroquine reduces fatality of critically ill patients with COVID-19. *Sci China Life Sci*. 2020:1–7. <https://doi.org/10.1007/s11427-020-1732-2>.
- [14] Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al. Remdesivir for the treatment of covid-19 - preliminary report. *N Engl J Med* 2020 May 22.
- [15] Chen Z, Hu J, Zhang Z, Jiang S, Han S, Yan D, Zhuang R, Hu B, Zhang Z. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. *medRxiv* 2020.03.22.20040758, <https://doi.org/10.1101/2020.03.22.20040758>; 2020.
- [16] Huang M, Tang T, Pang P, Li M, Ma R, Lu J, Shu J, You Y, Chen B, Liang J, Hong Z, Chen H, Kong L, Qin D, Pei D, Xia J, Jiang S, Shan H. Treating COVID-19 with chloroquine. *J Mol Cell Biol* 2020;12(4):322–5. <https://doi.org/10.1093/jmcb/mjaa014>.
- [17] Geleris J, Sun Y, Platt J, et al. Observational study of hydroxychloroquine in hospitalized patients with covid-19. *N Engl J Med* 2020;382(25):2411–8.
- [18] Liao X, Wang B, Kang Y. Novel coronavirus infection during the 2019-2020 epidemic: preparing intensive care units-the experience in Sichuan Province, China. *Intensive Care Med* 2020;46:357–60.
- [19] Armstrong N, Richez M, Raoult D, et al. Simultaneous UHPLC-UV analysis of hydroxychloroquine, minocycline and doxycycline from serum samples for the therapeutic drug monitoring of Q fever and Whipple's disease. *J Chromatogr B Analyt Technol Biomed Life Sci* 2017;1060:166–72.
- [20] Edouard S, Colson P, Melenotte C, et al. Evaluating the serological status of COVID-19 patients using an indirect immunofluorescent assay, France. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.05.20092064>.
- [21] <https://www.uptodate.com/contents/treatment-of-community-acquired-pneumonia-in-adults-who-require-hospitalization>.
- [22] Charlson M, Sztrowski TP, Peterson J, et al. Validation of a combined comorbidity index. *J Clin Epidemiol* 1994 Nov;47(11):1245–51.
- [23] Onder G, Rezza G, Brusaferro S. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *JAMA* 2020;23. <https://doi.org/10.1001/jama.2020.4683>. Published online March.
- [24] Zhang W, Qian BY. Making decisions to mitigate COVID-19 with limited knowledge. pii: S1473-3099 Lancet Infect Dis 2020 Apr 7;(20):30280–2. [https://doi.org/10.1016/S1473-3099\(20\)30280-2](https://doi.org/10.1016/S1473-3099(20)30280-2) [Epub ahead of print].
- [25] Décret n°2020-293 du 23 mars 2020, <https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000041746694&categorieLien=id>.

- [26] Couzin-Frankel J. The mystery of the pandemic's 'happy hypoxia. *Science* 2020; 368(6490):455–6. <https://doi.org/10.1126/science.368.6490.455>.
- [27] Davido B, Lansaman T, Bessis S, et al. Hydroxychloroquine plus azithromycin: a potential interest in reducing in-hospital morbidity due to COVID-19 pneumonia (HI-ZY-COVID)? medRxiv 2020. <https://doi.org/10.1101/2020.05.05.20088757>.
- [28] Mahévas M, Viet-Thi Tran, Roumier M, et al. Clinical efficacy of hydroxychloroquine in patients with covid-19 pneumonia who require oxygen: observational comparative study using routine care data. *BMJ* 2020;369.
- [29] Zeng QL, Yu ZJ, Gou JJ, et al. Effect of convalescent plasma therapy on viral shedding and survival in COVID-19 patients. *J Infect Dis* 2020;22(1):38–43.
- [30] Katzen J, Kohn R, Houk JL, et al. Early oseltamivir after hospital admission is associated with shortened hospitalization: a 5-year analysis of oseltamivir timing and clinical outcomes. *Clin Infect Dis* 2019 Jun 18;69(1):52–8.
- [31] Carlucci P, Ahuja T, Petrilli CM, et al. Hydroxychloroquine and azithromycin plus zinc vs hydroxychloroquine and azithromycin alone: outcomes in hospitalized COVID-19 patients. medRxiv 2020. <https://doi.org/10.1101/2020.05.02.20080036>. 05.02.20080036.
- [32] Rosenberg ES, Dufort EM, Udo T, et al. Association of treatment with hydroxychloroquine or azithromycin with in-hospital mortality in patients with COVID-19 in New York state. *JAMA* 2020;323(24):2493–502. May 11.
- [33] Middeldorp S, Coppens M, van Haaps TF, et al. Incidence of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemostasis* 2020 May 5 [Epub ahead of print].
- [34] Qin C, Zhou L, Hu Z, et al. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. *Clin Infect Dis* 2020 Mar 12. pii: ciaa248.
- [35] Zhang T, Sun LX, Feng RE. [Comparison of clinical and pathological features between severe acute respiratory syndrome and coronavirus disease 2019]. *0 Zhonghua Jiehe He Huxi Zazhi* 2020 Apr 3;43:E040 [Epub ahead of print].
- [36] Risch HA. Early outpatient treatment of symptomatic, high-risk covid-19 patients that should be ramped-up immediately as Key to the pandemic crisis. kwaa093 *Am J Epidemiol* 2020 May 27. <https://doi.org/10.1093/aje/kwaa093>. Online ahead of print.
- [37] Concato J, Shah N, Horwitz RI. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 2000 Jun 22;342(25): 1887–92. <https://doi.org/10.1056/NEJM200006223422507>.
- [38] Frieden TR. Evidence of health decision making-beyond randomized controlled trials. *N Engl J Med* 2017;377:465–75.
- [39] Gautret P, Raoult D. Nullane salus extra ecclesiam. *New Microbes New Infect* 2020. in press.